

It is time to move beyond 'body region silos' to manage musculoskeletal pain: five actions to change clinical practice

J P Caneiro ^{1,2}, Ewa M Roos ³, Christian J Barton,⁴
Kieran O'Sullivan ⁵, Peter Kent,^{6,7} Ivan Lin ⁸, Peter Choong,⁹
Kay M Crossley ⁴, Jan Hartvigsen,³ Anne Julia Smith,¹
Peter O'Sullivan¹

Current clinical research, education and practice commonly approaches musculoskeletal pain conditions in silos. A focus on body regions such as knee, hip, neck, shoulder and back pain as separate entities is manifest by region-specific clinical guidelines, conferences and working groups. Emerging evidence demonstrates that musculoskeletal pain disorders are frequently comorbid and share common biopsychosocial risk profiles for pain and disability.¹⁻⁵ There is broad consensus across clinical guidelines on the recommendations for best practice, irrespective of body region.³ We contend that a shift to focus on the person is needed. This best practice approach will encourage clinicians to (1) focus on patients' context and modifiable biopsychosocial factors that influence their pain and disability³; (2) use education to facilitate active management approaches (targeted exercise therapy, physical activity and healthy lifestyle habits) and reduce reliance on passive interventions; and (3) consider evidence-based surgical procedures only for those

with a clear indication and where guideline-based non-surgical approaches have been rigorously adhered to.

WE RECOMMEND FIVE ACTIONS TO MANAGE A PERSON WITH MUSCULOSKELETAL PAIN, IRRESPECTIVE OF BODY REGION

To adopt a person-centred active approach to treating musculoskeletal pain and disability, clinicians should:

1. Screen for biopsychosocial factors and health comorbidities

Clinicians need to communicate clearly with the patient to identify potential biopsychosocial drivers of pain and disability, which become therapeutic targets for their care.⁶⁻⁹ These include pain beliefs, emotional and coping responses to pain, social context, physical and lifestyle factors, as well as the presence of comorbidities. Clinicians should incorporate musculoskeletal screening tools, such as the Orebro Musculoskeletal Pain Questionnaire,¹⁰ to identify barriers to recovery and therapeutic targets for examination and care.

2. Embrace patient-centred communication

Clinicians should use open and reflective questioning to elicit the patient's understanding of factors, including (1) their pain experience ('Tell me your story'), (2) causation beliefs ('What do you think is the cause of your pain?'), (3) coping ('What do you do when pain increases?'), (4) impact ('Tell me how your symptoms affected your ability to engage with functional and physical activity'), (5) concerns ('Do your symptoms worry you?'), (6) beliefs regarding activity engagement ('Why do you think you shouldn't bend/lift/run?'), (7) social factors ('Tell me about your home/work/social life'), (8) goals ('Tell me about your goals') and (9) expectations

3. What do you think you need to achieve your goals?

This information can then guide an examination that explores the patient's concerns, functional limitations and physical capacity linked to their goals. Communication that privileges the patient's narrative results in patient-centred care and effective shared decision-making about potential risks and benefits of various interventions.^{5 6 9}

4. Educate beyond words using active learning approaches

Clinicians should embrace education as a central aspect of patient care in order to impart evidence-based information and to facilitate behavioural change. This involves the use of simple and clear language to dispel myths about pain, imaging findings and activity engagement.^{7 8 11-13} Education is an active process that is enhanced by writing down key information, using patient stories, web resources and learning aids (see online supplementary appendix). The formats used will vary according to patient's preferences and context. Behavioural learning, such as through exercise therapy, can be used to dispel unhelpful beliefs and build self-efficacy regarding the safety and benefits of movement and activity engagement. To evaluate learning, clinicians may ask patients to report 'take away' messages from each session.

5. Coach towards self-management

Clinicians should empower patients to engage in exercise, valued activities and a healthy lifestyle with confidence. Active management relieves pain and improves function across pain conditions and health comorbidities.^{14 15} This may involve group-based or individualised models of care. For some, this transition may be easy, while for others, this may reflect a gradual journey over an extended period of time. During this process, care is tailored to patients' individual needs and may include targeting (1) unhelpful cognitions (ie, developing a positive mindset and biopsychosocial understanding of their pain); (2) physical barriers to recovery (ie, load management and a graduated exercise programme of adequate dosage (intensity and frequency) addressing specific physical impairments such as weakness, mobility, movement avoidance and/or protective muscle guarding); and (3) lifestyle factors (ie, sleep, regular physical activity, relaxation, weight control, nutrition and positive social interactions). Clinicians should encourage patients to apply learnt strategies to daily life and

¹School of Physiotherapy and Exercise Science, Curtin University, Perth, Western Australia, Australia

²Body Logic Physiotherapy Clinic, Perth, Western Australia, Australia

³Center for Muscle and Joint Health, Department of Sports Science and Clinical Biomechanics, University of Southern Denmark, Odense, Denmark

⁴School of Allied Health, La Trobe University and Exercise Medicine Research Centre, Melbourne, Victoria, Australia

⁵University of Melbourne Department of Surgery, St Vincent's Hospital, Melbourne, New South Wales, Australia

⁶Aspetar Orthopaedic and Sports Medicine Hospital, Doha, Qatar

⁷School of Allied Health, University of Limerick, Limerick, Ireland

⁸Ageing Research Centre, University of Limerick, Limerick, Ireland

⁹Western Australian Centre for Rural Health, University of Western Australia, Geraldton, Western Australia, Australia

Correspondence to Dr J P Caneiro, Physiotherapy, School of Physiotherapy and Exercise Science, Curtin University, Bentley, WA 6102, Australia; j.p.caneiro@postgrad.curtin.edu.au

during periods of symptom exacerbation and distress. Booster sessions may be required during this process. Clinicians should teach patients how to self-monitor and progress their own programme. These strategies can help patients build self-efficacy and develop sustainable self-management strategies that are in line with their context, preferences and goals.

6. Address comorbid health factors

Clinicians should refer for co-care in the presence of comorbid mental and physical health complaints such as high levels of emotional distress, eating disorders (overeating and undereating) and type 2 diabetes. We contend that multidisciplinary care needs to be integrated, with consistent messages across the team to prevent care fragmentation and patient distress.

HEALTH SYSTEMS NEED TO SUPPORT CLINICIANS

Patient-centred care will optimise the value of healthcare provided. Shifting funding to support high-value evidence-based care options and educating society will be critical to enable this transition and will likely be cost-effective. Integrated cross-discipline clinical networks are required for effective co-care. We believe clinicians are ready to change, but they require the support of health systems and payers.

Twitter J P Caneiro @jpcaneiro, Ewa M Roos @ewa_roos, Christian J Barton @DrChrisBarton and Kieran O'Sullivan @kieranosull

Contributors JPC, POS, EMR and CJB developed the concept. JPC drafted the manuscript. All authors contributed to and approved the final version submitted.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests Several authors (JPC, CJB, KOS, KMC, and POS) are members of the editorial board of *British Journal of Sports Medicine*. Four authors (JPC, POS, KOS and CJB) deliver educational workshops on patient-centred care.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2018-100488>).



To cite Caneiro JP, Roos EM, Barton CJ, *et al.* *Br J Sports Med* Epub ahead of print: [please include Day Month Year]. doi:10.1136/bjsports-2018-100488

Accepted 16 September 2019

Br J Sports Med 2019;0:1–2.
doi:10.1136/bjsports-2018-100488

ORCID iDs

J P Caneiro <http://orcid.org/0000-0001-5555-4412>
Ewa M Roos <http://orcid.org/0000-0001-5425-2199>
Kieran O'Sullivan <http://orcid.org/0000-0002-7137-3125>
Ivan Lin <http://orcid.org/0000-0001-6901-2569>
Kay M Crossley <http://orcid.org/0000-0001-5892-129X>

REFERENCES

- Kittelson AJ, George SZ, Maluf KS, *et al.* Future directions in painful knee osteoarthritis: harnessing complexity in a heterogeneous population. *Phys Ther* 2014;94:422–32.
- Hannibal KE, Bishop MD. Chronic stress, cortisol dysfunction, and pain: a psychoneuroendocrine rationale for stress management in pain rehabilitation. *Phys Ther* 2014;94:1816–25.
- Lin I, Wiles L, Waller R, *et al.* What does best practice care for musculoskeletal pain look like? eleven consistent recommendations from high-quality clinical practice guidelines: systematic review. *Br J Sports Med* 2019;bjsports-2018-099878.
- Hartvigsen J, Hancock MJ, Kongsted A, *et al.* What low back pain is and why we need to pay attention. *Lancet* 2018;391:2356–67.
- Lewis J, O'Sullivan P. Is it time to reframe how we care for people with non-traumatic musculoskeletal pain? *Br J Sports Med* 2018;52:1543–4.
- Synnott A, O'Keeffe M, Bunzli S, *et al.* Physiotherapists report improved understanding of and attitude toward the cognitive, psychological and social dimensions of chronic low back pain after cognitive functional therapy training: a qualitative study. *J Physiother* 2016;62:215–21.
- MacKay C, Hawker GA, Jaglal SB. Qualitative study exploring the factors influencing physical therapy management of early knee osteoarthritis in Canada. *BMJ Open* 2018;8:e023457.
- Cowell I, O'Sullivan P, O'Sullivan K, *et al.* Perceptions of physiotherapists towards the management of non-specific chronic low back pain from a biopsychosocial perspective: a qualitative study. *Musculoskelet Sci Pract* 2018;38:113–9.
- Edmond SN, Keefe FJ. Validating pain communication: current state of the science. *Pain* 2015;156:215–9.
- Linton SJ, Nicholas M, MacDonald S. Development of a short form of the Örebro musculoskeletal pain screening questionnaire. *Spine* 2011;36:1891–5.
- Jenkins HJ, Hancock MJ, Maher CG, *et al.* Understanding patient beliefs regarding the use of imaging in the management of low back pain. *European Journal of Pain* 2016;20:573–80.
- O'Keeffe M, O'Sullivan PB, O'Sullivan K. Education can 'change the world': Can clinical education change the trajectory of individuals with back pain? *Br J Sports Med* 2019;bjsports-2018-100190.
- Buchbinder R, Staples M, Jolley D. Doctors with a special interest in back pain have poorer knowledge about how to treat back pain. *Spine* 2009;34:1218–26. discussion 27.
- Skou ST, Pedersen BK, Abbott JH, *et al.* Physical activity and exercise therapy benefit more than just symptoms and impairments in people with hip and knee osteoarthritis. *J Orthop Sports Phys Ther* 2018;48:439–47.
- Pedersen BK, Saltin B. Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports* 2015;25:1–72.